

# Introduction to the General Linear Model

Consider the General Linear Model

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon},$$

where ...

$y$  is an  $n \times 1$  random vector of responses that can be observed.

The values in  $y$  are values of the response variable.

$X$  is an  $n \times p$  matrix of known constants.

Each column of  $X$  contains the values for an explanatory variable that is also known as a predictor variable or a regressor variable in the context of multiple regression.

The matrix  $X$  is sometimes referred to as the design matrix.

$\beta$  is an unknown parameter vector in  $\mathbb{R}^p$ .

We are often interested in estimating a LC of the elements in  $\beta$  ( $c'\beta$ ) or multiple LCs ( $C\beta$ ) for some known  $c$  or  $C$ .

$\varepsilon$  is an  $n \times 1$  random vector that cannot be observed.

The values in  $\varepsilon$  are called errors. Initially, we assume only  $E(\varepsilon) = \mathbf{0}$ .

Note that

$$\begin{aligned} E(\boldsymbol{\varepsilon}) = \mathbf{0} &\Rightarrow E(\mathbf{y}) = E(\mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}) \\ &= \mathbf{X}\boldsymbol{\beta} + E(\boldsymbol{\varepsilon}) \\ &= \mathbf{X}\boldsymbol{\beta} \in \mathcal{C}(\mathbf{X}). \end{aligned}$$

Thus, this general linear model simply says that  $\mathbf{y}$  is a random vector with mean  $E(\mathbf{y})$  in the column space of  $X$ .



## Example:

Suppose 5 hogs are fed diet 1 for three weeks. Let  $y_i$  be the weight gain of the  $i^{\text{th}}$  hog for  $i = 1, \dots, 5$ .

Suppose that we assume  $E(y_i) = \mu \forall i = 1, \dots, 5$ . Then

$$\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \mu + \begin{bmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \varepsilon_3 \\ \varepsilon_4 \\ \varepsilon_5 \end{bmatrix}$$

where  $E(\varepsilon_i) = 0 \forall i = 1, \dots, 5$ .

## Example:

Suppose 10 hogs are randomly divided into two groups of 5 hogs each. Group 1 is fed diet 1 for three weeks, group 2 is fed diet 2 for three weeks.

Let  $y_{ij}$  denote the weight gain of the  $j^{\text{th}}$  hog in the  $i^{\text{th}}$  diet group ( $i = 1, 2; j = 1, \dots, 5$ ).

If we assume that  $E(y_{ij}) = \mu_i$  for  $i = 1, 2; j = 1, \dots, 5$ , then

$$\begin{bmatrix} y_{11} \\ y_{12} \\ y_{13} \\ y_{14} \\ y_{15} \\ y_{21} \\ y_{22} \\ y_{23} \\ y_{24} \\ y_{25} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 0 \\ 1 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix} + \begin{bmatrix} \varepsilon_{11} \\ \varepsilon_{12} \\ \varepsilon_{13} \\ \varepsilon_{14} \\ \varepsilon_{15} \\ \varepsilon_{21} \\ \varepsilon_{22} \\ \varepsilon_{23} \\ \varepsilon_{24} \\ \varepsilon_{25} \end{bmatrix}$$

where  $E(\varepsilon_{ij}) = 0 \forall i = 1, 2; j = 1, \dots, 5$ .

Alternatively, we could assume  $E(y_{ij}) = \mu + \tau_i \forall i = 1, 2; j = 1, \dots, 5$ .

Then the design matrix and parameter vector become ...

$$\begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} \mu \\ \tau_1 \\ \tau_2 \end{bmatrix} = \begin{bmatrix} \mu + \tau_1 \\ \mu + \tau_1 \\ \mu + \tau_1 \\ \mu + \tau_1 \\ \mu + \tau_1 \\ \mu + \tau_2 \\ \mu + \tau_2 \\ \mu + \tau_2 \\ \mu + \tau_2 \\ \mu + \tau_2 \end{bmatrix} .$$

These models are equivalent because both design matrices have the same column space:

$$\left\{ \begin{bmatrix} a \\ a \\ a \\ a \\ a \\ b \\ b \\ b \\ b \\ b \end{bmatrix} : a, b \in \mathbb{R} \right\} .$$



## Example:

10 steer carcasses were assigned to be measured for pH at one of five times after slaughter. Data are as follows (Schwenke & Milliken(1991), Biometrics, 47, 563-573.)

Steer	Hours after Slaughter	pH
1	1	7.02
2	1	6.93
3	2	6.42
4	2	6.51
5	3	6.07
6	3	5.99
7	4	5.59
8	4	5.80
9	5	5.51
10	5	5.36

$\forall i = 1, \dots, 10$ , let

$x_i$  = measurement time (hours after slaughter) for steer  $i$

$y_i$  = pH for steer  $i$ .

Suppose  $y_i = \beta_0 + \beta_1 \log(x_i) + \varepsilon_i$  where  $E(\varepsilon_i) = 0 \forall i = 1, \dots, 10$ .

Determine  $\mathbf{y}$ ,  $\mathbf{X}$ , and  $\boldsymbol{\beta}$ .

Chapter 1 of our text contains many more examples. Please read the entire chapter.