

William Q. Meeker and Luis A. Escobar
Iowa State University and Louisiana State University


Copyright 1998-2008 W. Q. Meeker and L. A. Escobar.
Based on the authors' text *Statistical Methods for Reliability Data*, John Wiley & Sons Inc. 1998.

July 30, 2009
8h 3min

Bootstrap Sampling and Bootstrap Confidence Intervals

- Instead of assuming $Z_{\hat{\mu}} = (\hat{\mu} - \mu) / \widehat{SE}_{\hat{\mu}} \sim \text{NOR}(0, 1)$, use Monte Carlo simulation to approximate the distribution of $Z_{\hat{\mu}}$.
 - Simulate $B = 4000$ values of $Z_{\hat{\mu}^*} = (\hat{\mu}^* - \hat{\mu}) / \widehat{SE}_{\hat{\mu}^*}$.
 - Some bootstrap approximations:
 - ▶ $Z_{\hat{\mu}} \sim Z_{\hat{\mu}^*}$
 - ▶ $Z_{\log(\hat{\sigma})} \sim Z_{\log(\hat{\sigma}^*)}$
 - ▶ $Z_{\logit[\hat{F}(t)]} \sim Z_{\logit[\hat{F}^*(t)]}$
- when computing confidence intervals for μ , σ , and F .


A Simple Bootstrap Re-Sampling Method




Bootstrap Confidence Intervals
Chapter 9 Objectives

- Explain basic ideas behind the use of computer simulation to obtain bootstrap confidence intervals.
- Explain different methods for generating bootstrap samples.
- Obtain and interpret simulation-based pointwise parametric bootstrap confidence intervals.


A Simple Parametric Bootstrap Sampling Method



Scatterplot of 1,000 (Out of $B = 10,000$) Bootstrap Estimates $\hat{\mu}^*$ and $\hat{\sigma}^*$ for Shock Absorber



Weibull Plot of $F(t; \hat{\mu}, \hat{\sigma})$ from the Original Sample (dark line) and 50 (Out of $B=10,000$) $F(t; \hat{\mu}^*, \hat{\sigma}^*)$ Computed from Bootstrap Samples for the Shock Absorber



9 - 7

Bootstrap Confidence Interval for μ

- With complete data or Type II censoring,

$$Z_{t_j^*} = \frac{\hat{\mu}_j^* - \hat{\mu}}{\widehat{SE}_{\hat{\mu}_j^*}}$$

has a distribution that does not depend on any unknown parameters. Such a quantity is called a **pivotal** quantity.

- By the definition of quantiles, then

$$\Pr \left(z_{t_j^*}^{(1-\alpha/2)} < Z_{t_j^*} \leq z_{t_j^*}^{(\alpha/2)} \right) = 1 - \alpha$$


- Simple algebra shows that

$$[\underline{\mu}, \bar{\mu}] = [\hat{\mu} - z_{t_j^*}^{(1-\alpha/2)} \widehat{SE}_{\hat{\mu}}, \hat{\mu} + z_{t_j^*}^{(\alpha/2)} \widehat{SE}_{\hat{\mu}}]$$

provides an exact 95% confidence interval for μ . With other kinds of censoring, the interval is, in general, only **approximate**.

9 - 8

Bootstrap Distributions of Weibull $\hat{\mu}^*$ and $Z_{t_j^*}$ Based on $B=10,000$ Bootstrap Samples for the Shock Absorber



9 - 9

Bootstrap Confidence Interval for σ

- With complete data or Type II censoring,

$$Z_{\log(\hat{\sigma}^*)} = \frac{\log(\hat{\sigma}^*) - \log(\hat{\sigma})}{\widehat{SE}_{\log(\hat{\sigma}^*)}}$$

has a distribution that does not depend on any unknown parameters. Such a quantity is called a **pivotal** quantity.

- By the definition of quantiles, then

$$\Pr \left(z_{\log(\hat{\sigma}^*)}^{(1-\alpha/2)} < Z_{\log(\hat{\sigma}^*)} \leq z_{\log(\hat{\sigma}^*)}^{(\alpha/2)} \right) = 1 - \alpha$$

- Simple algebra shows that


$$[\underline{\hat{\sigma}}, \bar{\hat{\sigma}}] = [\hat{\sigma}/\hat{w}, \hat{\sigma}/\hat{w}]$$

provides an exact 95% confidence interval for σ , where $\hat{w} =$

$\exp \left[z_{\log(\hat{\sigma}^*)}^{(1-\alpha/2)} \widehat{SE}_{\log(\hat{\sigma})} \right]$ and $\hat{w} = \exp \left[z_{\log(\hat{\sigma}^*)}^{(\alpha/2)} \widehat{SE}_{\log(\hat{\sigma})} \right]$ With other kinds of censoring, the interval is, in general, only **approximate**.

9 - 10

Bootstrap Distributions of $\hat{\sigma}^*$, $Z_{\hat{\sigma}^*}$, and $Z_{\log(\hat{\sigma}^*)}$ Based on $B=10,000$ Bootstrap Samples



9 - 11

Bootstrap Confidence Interval for $F(t_e)$

- With complete data or Type II censoring [using $F = F(t_e)$],

$$Z_{\logit(\hat{F}^*)} = \frac{\logit(\hat{F}^*) - \logit(\hat{F})}{\widehat{SE}_{\logit(\hat{F}^*)}}$$

has a distribution that does not depend on any unknown parameters. Such a quantity is called a **pivotal** quantity.

- By the definition of quantiles, then

$$\Pr \left(z_{\logit(\hat{F}^*)}^{(1-\alpha/2)} < Z_{\logit(\hat{F}^*)} \leq z_{\logit(\hat{F}^*)}^{(\alpha/2)} \right) = 1 - \alpha$$


- Simple algebra shows that

$$[\underline{F}, \bar{F}] = \left[\frac{\hat{F}}{\hat{F} + (1 - \hat{F}) \times \hat{w}}, \frac{\hat{F}}{\hat{F} + (1 - \hat{F}) \times \hat{w}} \right]$$

where provides an exact 95% confidence interval for F , where $\hat{w} = \exp \left[z_{\logit(\hat{F}^*)}^{(1-\alpha/2)} \widehat{SE}_{\logit(\hat{F})} \right]$ and $\hat{w} = \exp \left[z_{\logit(\hat{F}^*)}^{(\alpha/2)} \widehat{SE}_{\logit(\hat{F})} \right]$ With other kinds of censoring, the interval is, in general, only **approximate**.

9 - 12

Bootstrap Distributions of $\hat{F}(t_e)^*$, $Z_{\hat{F}(t_e)^*}$, and $Z_{\log[\hat{F}(t_e)^*]}$ for $t_e=10,000$ km Based on $B=10,000$ Bootstrap Samples



9 - 13

Bootstrap Confidence Interval for t_p

- With complete data or Type II censoring, $Z_{\log(\hat{t}_p^*)} = \frac{\log(\hat{t}_p^*) - \log(\hat{t}_p)}{\widehat{SE}_{\log(\hat{t}_p^*)}}$ has a distribution that does not depend on any unknown parameters. Such a quantity is called a **pivotal** quantity.
- By the definition of quantiles, then

$$\Pr \left(z_{\log(\hat{t}_p^*)}(\alpha/2) < Z_{\log(\hat{t}_p^*)} \leq z_{\log(\hat{t}_p^*)}(1-\alpha/2) \right) = 1 - \alpha$$


- Simple algebra shows that

$$[\hat{t}_p, \hat{t}_p] = [\hat{t}_p/\hat{w}, \hat{t}_p/\hat{w}]$$

provides an exact 95% confidence interval for t_p , where $\hat{w} = \exp \left[z_{\log(\hat{t}_p^*)}(1-\alpha/2) \widehat{SE}_{\log(\hat{t}_p)} \right]$ and $\hat{w} = \exp \left[z_{\log(\hat{t}_p^*)}(\alpha/2) \widehat{SE}_{\log(\hat{t}_p)} \right]$. With other kinds of censoring, the interval is, in general, only **approximate**.

9 - 14

Bootstrap Distributions of \hat{t}_p , $Z_{\hat{t}_p}$, and $Z_{\log[\hat{t}_p]}$ for $t_e=10,000$ km Based on $B=10,000$ Bootstrap Samples



9 - 15